

Already in EDAN

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1652KHG

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the
IPC reform
NEWS 4 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/
USPAT2
NEWS 5 JAN 13 IPC 8 searching in IFIPAT, IFIUDB, and IFICDB
NEWS 6 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to
INPADOC
NEWS 7 JAN 17 Pre-1988 INPI data added to MARPAT
NEWS 8 JAN 17 IPC 8 in the WPI family of databases including WPIFV
NEWS 9 JAN 30 Saved answer limit increased
NEWS 10 JAN 31 Monthly current-awareness alert (SDI) frequency
added to TULSA
NEWS 11 FEB 21 STN AnaVist, Version 1.1, lets you share your STN AnaVist
visualization results
NEWS 12 FEB 22 Status of current WO (PCT) information on STN
NEWS 13 FEB 22 The IPC thesaurus added to additional patent databases on STN
NEWS 14 FEB 22 Updates in EPFULL; IPC 8 enhancements added
NEWS 15 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 16 FEB 28 MEDLINE/LMEDLINE reload improves functionality
NEWS 17 FEB 28 TOXCENTER reloaded with enhancements
NEWS 18 FEB 28 REGISTRY/ZREGISTRY enhanced with more experimental spectral
property data
NEWS 19 MAR 01 INSPEC reloaded and enhanced
NEWS 20 MAR 03 Updates in PATDPA; addition of IPC 8 data without attributes
NEWS 21 MAR 08 X.25 communication option no longer available after June 2006

NEWS EXPRESS FEBRUARY 15 CURRENT VERSION FOR WINDOWS IS V8.01a,
CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005.
V8.0 AND V8.01 USERS CAN OBTAIN THE UPGRADE TO V8.01a AT
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 15:17:17 ON 16 MAR 2006

=> File HCAPLUS

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'HCAPLUS' ENTERED AT 15:17:40 ON 16 MAR 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Mar 2006 VOL 144 ISS 12

FILE LAST UPDATED: 15 Mar 2006 (20060315/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s (gamma-type pyridoxal phosphate enzymes)

805911 GAMMA

4916 GAMMAS

806071 GAMMA

(GAMMA OR GAMMAS)

1658769 TYPE

574761 TYPES

2103189 TYPE

(TYPE OR TYPES)

13514 PYRIDOXAL

54 PYRIDOXALS

13520 PYRIDOXAL

(PYRIDOXAL OR PYRIDOXALS)

537851 PHOSPHATE

125764 PHOSPHATES

586208 PHOSPHATE

(PHOSPHATE OR PHOSPHATES)

442972 ENZYMES

L1 0 (GAMMA-TYPE PYRIDOXAL PHOSPHATE ENZYMES)

(GAMMA(W) TYPE(W) PYRIDOXAL(W) PHOSPHATE(W) ENZYMES)

=> s (pyridoxal phosphate enzymes)

13514 PYRIDOXAL

54 PYRIDOXALS

13520 PYRIDOXAL

(PYRIDOXAL OR PYRIDOXALS)

537851 PHOSPHATE

125764 PHOSPHATES

586208 PHOSPHATE

(PHOSPHATE OR PHOSPHATES)

442972 ENZYMES

L2 82 (PYRIDOXAL PHOSPHATE ENZYMES)
 (PYRIDOXAL(W) PHOSPHATE(W) ENZYMES)

=> s L2 and uncinatum
 263 UNCINATUM

L3 0 L2 AND UNCINATUM

=> s L2 and alkaloid
 53415 ALKALOID
 67894 ALKALOIDS
 81309 ALKALOID
 (ALKALOID OR ALKALOIDS)

L4 0 L2 AND ALKALOID

=> s neotyphodium and L2
 192 NEOTYPHODIUM

L5 0 NEOTYPHODIUM AND L2

=> s l2 and fungal
 48730 FUNGAL
 8 FUNGALS
 48734 FUNGAL
 (FUNGAL OR FUNGALS)

L6 0 L2 AND FUNGAL

=> s Spiering
 7 SPIERING
 2 SPIERINGS
L7 9 SPIERING
 (SPIERING OR SPIERINGS)

=> d 1-9 abs,so,ti

L7 ANSWER 1 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

AB Single-mol. fluorescence resonance energy transfer and functional assays have been used to study the initiation and regulation of the bacteriophage T4 DNA replication system. Previous work has demonstrated that a complex of the helicase loading protein (gp59) and the DNA polymerase (gp43) on forked DNA totally inhibits the polymerase and exonuclease activities of gp43 by a mol. locking mechanism (Xi, J., Zhuang, Z., Zhang, Z., Selzer, T., Spiering, M. M., Hammes, G. G., and Benkovic, S. J. (2005) Biochem. 44, 2305-2318). We now show that this complex is "unlocked" by the addition of the helicase (gp41) with restoration of the DNA polymerase activity. Gp59 retains its ability to load the helicase while forming a gp59-gp43 complex at a DNA fork in the presence of the single-stranded DNA binding protein (gp32). Upon the addition of gp41 and MgATP, gp59 dissociates from the complex, and the DNA-bound gp41 is capable of recruiting the primase (gp61) to form a functional primosome and, subsequently, a fully active replisome. Functional assays of leading- and lagging-strand synthesis on an active replication fork show that the absence of gp59 has no effect on the coupling of leading- and lagging-strand synthesis or on the size of the Okazaki DNA fragments. We conclude that gp59 acts in a manner similar to the clamp loader to ensure proper assembly of the replisome and does not remain as a replisome component during active replication.

SO Biochemistry (2005), 44(21), 7747-7756
 CODEN: BICHAW; ISSN: 0006-2960

TI Interaction between the T4 Helicase Loading Protein (gp59) and the DNA Polymerase (gp43): Unlocking of the gp59-gp43-DNA Complex to Initiate Assembly of A Fully Functional Replisome

L7 ANSWER 2 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

AB The name of the sixth author, Barry A. Spiering, was misspelled.

- SO Medicine & Science in Sports & Exercise (2004), 36(2), 349
CODEN: MSPEDA; ISSN: 0195-9131
- TI Effect of muscle oxygenation during resistance exercise on anabolic hormone response. [Erratum to document cited in CA140:193472]
- L7 ANSWER 3 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN
- AB The Monte Carlo entropic sampling method previously presented is adapted here to an Ising-like system with short- and long-range interactions. Such model is suited to spin crossover solids where the long interaction is due to elastic coupling mediated by the lattice, while the short-range interaction originates from the bonding between the spin crossover units [J. Linares, H. Spiering, F. Varret, Eur. J. Phys. B 10(1999) 271]. Taking into account the different degeneracies g_{HS} for high-spin (HS) and g_{LS} for low-spin (LS) states, the Ising Hamiltonian associated with fictitious spins is written: $H = -h \sum_i \sigma_i - J \sum_{\langle i,j \rangle} \sigma_i \sigma_j$ with $h = -\Delta + k_B T \ln(g_{HS} g_{LS}^2 + G \langle \sigma \rangle)$ where J and G are the short- and long-range interactions, resp., and Δ the energy gap of ligand field such that the LS state is the ground state. The numerical method was tested successfully by comparison to the exact solution for a 1-dimensional system: $[\text{Fe}(\text{Htrz})_2(\text{trz})](\text{BF}_4)_2$ []. The authors describe here the results obtained for 2-dimensional systems, and show that the squareness of the thermal hysteresis loop, associated with the spin-transition, can be correlated to the strength of short-range interactions.
- SO Polyhedron (2003), 22(14-17), 2453-2456
CODEN: PLYHDE; ISSN: 0277-5387
- TI Monte Carlo entropic sampling applied to spin crossover solids: the squareness of the thermal hysteresis loop
- L7 ANSWER 4 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN
- AB The phenomenon of thermal spin crossover in iron(II) complexes has been subject to vigorous research activities in recent years, with the ultimate goal of eventual tech. applications in switching and display devices. The changes of spin states with accompanying drastic changes in magnetic and optical properties upon variation of temperature or application of pressure or irradiation with light has been explored extensively, particularly in view of the cooperative interactions playing a decisive role in the solid state [1]. Remarkable progress has been made in the design and preparation of new spin crossover materials, new photophys. effects have been observed, studies of the spin transition behavior under pressure have yielded astonishing new results, structural investigations at variable temperature and after irradiation with light have permitted deeper insight into the lattice events accompanying the mol. switching processes. Some of the new results will be highlighted in this report. [1/ P. Gutlich, A. Hauser, H. Spiering, Angew. Chemical Int. Ed. Engl. 1994, 33, 2024-2054.
- SO Book of Abstracts, 219th ACS National Meeting, San Francisco, CA, March 26-30, 2000 (2000), INOR-176 Publisher: American Chemical Society, Washington, D. C.
CODEN: 69CLAC
- TI New highlights in spin crossover studies.
- L7 ANSWER 5 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN
- AB In theory, the hydration thermodyn. approach used for control of glass production in the Defense Waste Processing Facility should be applicable to virtually any situation in which glass is allowed to react with an aqueous medium. In this paper, the general approach has been modified for application to glass etching by aqueous HF. Using measurements made by G. Spierings (1991), who etched a wide variety of silicate and borosilicate compns. in 2.9 M HF, excellent agreement is found between the theory and the exptl. measurements.
- SO Ceramic Transactions (1998), 87(Environmental Issues and Waste Management Technologies in the Ceramic and Nuclear Industries III), 489-497

CODEN: CETREW; ISSN: 1042-1122

TI Thermodynamic approach to glass etching by HF

L7 ANSWER 6 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

AB An Ising-type model for spin conversion, explicitly accounting for intramol. vibrations was studied. Each two level system is associated with p harmonic oscillators having two possible frequencies $\omega(i)_{LS}$, $\omega(i)_{HS}$. The major advantage of this model is that it provides an excellent agreement with both the conversion curve and calorimetric data, in particular the entropy change upon spin conversion. With the help of Arrhenius plots, the predictions of the model are accurately compared to the literature exptl. data on the spin equilibrium curves. A very accurate Moessbauer study by Jacobi, Spiering and Gutlich, provides evidence for a small effect typical for vibrations. A novel example is given where the conversion curve is essentially monitored by vibrations; this originates from the extremely small value of the energy gap between HS and LS electro-vibrational ground states. However, in most cases, as a 1st approach, the low frequency approximation of the model can be used. Then the model reduces to a simple two-level model with addnl. degeneracies implicitly accounting for intramol. vibrations.

SO Journal de Physique I (1995), 5(6), 747-60

CODEN: JPGCE8; ISSN: 1155-4304

TI A simple Ising-like model for spin conversion including molecular vibrations

L7 ANSWER 7 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

AB The thermodyn. and kinetics of the HS .dblharw. LS equilibrium in the iron(II) spin-crossover complexes $[\text{Fe}(\text{2-pic})_3](\text{PF}_6)_2$ (2-pic = 2-aminomethylpyridine), and $[\text{Fe}(\text{6-mepy})_3\text{tren}](\text{PF}_6)_2$ (6-mepy = tris[4-[(6-methyl)-2-pyridyl]-3-aza-3-butenyl]amine) were studied by pressure dependent Moessbauer spectroscopy. The rate consts. for the HS .dblharw. LS intersystem crossing are determined by lineshape anal. of Moessbauer spectra. The evaluation and the essential features of quadrupole-quadrupole relaxation spectra are discussed. From thermodyn. anal. of the HS fractions the intermol. interaction consts. Γ and the volume changes of the lattice ΔV_c associated with the spin-transition are derived. In $[\text{Fe}(\text{2-pic})_3](\text{PF}_6)_2$, which shows a very gradual spin-transition, Γ vanishes and a small $\Delta V_c = 8\text{\AA}^3$ is found. For $[\text{Fe}(\text{6-mepy})_3\text{tren}](\text{PF}_6)_2$, which shows a steep spin-transition, $\Gamma = 197\text{ cm}^{-1}$ and $\Delta V_c = 25\text{\AA}^3$ are derived. The activation energy of the exothermic HS \rightarrow LS conversion is pressure independent for the pic-complex, but decreases with increasing pressure for the tren-complex. This is a consequence of different activation vols. ΔV_{HL}^\ddagger . For $[\text{Fe}(\text{6-mepy})_3\text{tren}](\text{PF}_6)_2$ $\Delta V_{HL}^\ddagger = -22\text{\AA}^3$ was estimated. The thermodyn. and kinetic results are discussed within the lattice expansion model of H Spiering et al. (1982) who treated the lattice influence on the spin state conversion.

SO Journal of Physics and Chemistry of Solids (1989), 50(6), 587-97

CODEN: JPCSAW; ISSN: 0022-3697

TI Thermodynamics and kinetics of spin state conversion processes studied by pressure dependent Moessbauer spectroscopy

L7 ANSWER 8 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN

AB A polemic with C. Srikanthappa, C. Raith, and B. Spiering (ibid., 1985, 26, 249-72). At the Kabbaldurga area (India), H₂O-fluid inclusions dominate the amphibolite-facies rocks. Since no partial melting took place, the structural and metamorphic setting of the Ponmudi charnockites are quite different from the Kabbaldurga occurrence, where partial melting was important. A body the size of the Closepet granite is unlikely to have originated by K metasomatism. The Closepet-type granites did not intrude only after charnockitization of gneisses. Fluid conditions must be studied to determine whether the influx of CO₂ was oxidizing or reducing.

SO Journal of the Geological Society of India (1988), 31(1), 166-71

CODEN: JGSIAJ; ISSN: 0016-7622

- TI Progressive charnockitization of a leptynite-khondalite suite from Kerala, India. Evidence for formation of charnockites through decrease in fluid pressure? Comment
- L7 ANSWER 9 OF 9 HCAPLUS COPYRIGHT 2006 ACS on STN
- AB A polemic. The results reported by Spierings and Stein (ibid. 1975, 6, 265) on the effect of Na₂O on hydration of C3A [12042-78-3] indicate that the authors are not aware of a series of (5) papers published by P. Gupta et. al.
- SO Cement and Concrete Research (1977), 7(1), 105
CODEN: CENRAI; ISSN: 0008-8846
- TI A discussion of the paper "The influence of sodium oxide on the hydration of C3A. I. Paste hydration", by G. A. C. M. Spierings and H. N. Stein